

RESEARCH

Open Access



# Expression patterns of *TRa* and *CRABP2* genes in Chinese cashmere goat skin during prenatal development

Tao Zhong<sup>1†</sup>, Wei Zhao<sup>1†</sup>, Zhongqiang Zhou<sup>1</sup>, Li Li<sup>1</sup>, Linjie Wang<sup>1</sup>, Hua Li<sup>2\*</sup> and Hongping Zhang<sup>1\*</sup>

## Abstract

**Background:** The physiologic characteristics of the cashmere trait and many of the differentially expressed genes relevant to hair cycling have been extensively studied, whereas genes involved in the prenatal development of hair follicles have been poorly investigated in cashmere goats. The aim of this study, therefore, was to quantify the time-course changes in the expressions of *TRa* and *CRABP2* genes in the fetal skin of Chinese cashmere goats at the multiple embryonic days (E70, E75, E80, E90, E100, E120 and E130) using real-time quantitative PCR (RT-qPCR).

**Results:** RT-qPCR showed that *TRa* was expressed at E70 with relatively high level and then slightly decreased (E75, E80, and E90). The highest expression of *TRa* mRNA was revealed at E130 ( $P > 0.05$ ). The expression pattern of *CRABP2* mRNA showed an 'up-down-up' trend, which revealed a significantly highest expression at E75 ( $P < 0.05$ ) and was down-regulated during E80 to E120 ( $P < 0.05$ ) and mildly increased at E130, subsequently.

**Conclusion:** This study demonstrated that *TRa* and *CRABP2* genes expressed in different levels during prenatal development of cashmere. The present study will be helpful to provide the comprehensive understanding of *TRa* and *CRABP2* genes expressions during cashmere formation and lay the ground for further studies on their roles in regulation of cashmere growth in goats.

**Keywords:** Cashmere goat, *TRa*, *CRABP2*, Skin, Expression

## Background

The Inner Mongolian cashmere goat is a Chinese indigenous breed characterized as a double-coated species. The outer coat consists of coarse guard hairs and the undercoat is the soft and precious cashmere. Two kinds of hair follicles which known as primary hair follicles and secondary hair follicles existed in the skin of the Inner Mongolian cashmere goat. Cashmere, which is derived from the secondary hair follicles, has smaller diameters than wool fibers produced by the primary hair follicles. Primary hair follicles and secondary hair follicles form at different periods and play different roles in the development of hair. In mice, the primary hair follicles arose *in utero* from

embryonic day (E) 12.5 and the secondary follicles started to develop until E17 [1, 2]. In goat embryos, the precursor primary follicles were observed in head, neck, shoulder, and belly at E45. The hair follicles gradually formed during 55E to 65E and developed into the mature primary follicles at E135 [3]. The morphogenesis and development of the secondary follicles were similar to those of the primary follicles. The secondary follicles grew from E65 to E75 and then extended to skin surface. The complete structure of the secondary follicle was formed at E135 in the embryos of Chinese cashmere goats [4]. Furthermore, the periodic growth of the secondary follicles also presented in a breed-specific manner [5]. All primary follicles but few secondary follicles were mature at birth and the number of secondary follicles increased 10-fold in the 57 days after birth. The number of primary follicles showed a tendency to decline between 57 and 107 days of age in Australian cashmere goats [6]. Like the hair follicle cycling in other mammals, the growth

\* Correspondence: 121183050@qq.com; zhp@sicau.edu.cn

<sup>†</sup>Equal contributors

<sup>2</sup>School of Life Science, Foshan University, Foshan 528000, P. R. China

<sup>1</sup>Farm Animal Genetic Resources Exploration and Innovation Key Laboratory of Sichuan Province, College of Animal Science and Technology, Sichuan Agricultural University, Chengdu, Sichuan 611130, P. R. China

of cashmere in goats was also tightly programmed by the three synchronized interchanging stages, anagen (growth phase), catagen (regression phase) and telogen (resting phase) throughout postnatal life [7, 8].

Some genes involved in the growth and development of hair follicles in cashmere goats have been identified, such as *KAPs* [9, 10], *BMP* [11], *Prolactin* [12], and *Keratin* [13, 14]. Furthermore, in mammals, the thyroid hormones (TH) have multi-functions in many important physiological processes including the normal growth, development, differentiation and metabolism. Recently, new insights into TH biological function have been obtained from animal studies involving in epidermis, dermis and hair cycling including anagen prolongation and stimulation of both hair matrix keratinocyte proliferation and hair pigmentation [15, 16]. The metabolism of TH is related to deiodinase, which is also regulated cashmere growth by altering its activity in skin tissue [17, 18]. TH action could be mediated through the thyroid hormone receptor (TR), which is part of the nuclear hormone receptor superfamily and bound to TH in three patterns identified in skin [19, 20]. TR interacts with the hairless gene product, a transcription factor required for hair growth. TR has been detected in epidermal keratinocytes, skin fibroblasts and a number of cell types that made up the hair follicles. In addition, the retinoic acid (RA) is essential for the development and maintenance of hair cycling [21]. The cellular RA-binding protein type II (CRABPII) is involved in RA synthesis pathway, which could shuttle RA to its receptor in nucleus and increase its

transcriptional efficiency [22]. Therefore, the dynamic expression of *CRABPII* mRNA could affect the concentration of RA, which acts on the formation of cashmere though regulating sebaceous gland. During hair cycle, the expression pattern of the RA synthesis and signaling including *Crbp*, *Dhrs9*, *Aldh1a1*, *Aldh1a2*, *Aldh1a3* and *Crabp2* defined in rodents only [23].

Based on the genetic studies in humans and rodents, *TR $\alpha$*  and *CRABPII* acted important roles in driving the progression of the hair cycle. We postulate that these two genes might have functions during cashmere formation in goat. So in this study, we described the characteristics of *TR $\alpha$*  and *CRABPII* genes in the Inner Mongolian cashmere goat and identified their expression patterns in skin tissue during the middle late embryonic stages (E70 to E130).

## Materials and methods

### Animal and skin tissue preparation

The Inner Mongolia cashmere goat is a traditional outstanding breed, which is famous for its excellent cashmere performance and strong adaptation to the semi-desert and desert steppes. The tested individuals were selected from the Aerbasi White Cashmere Goat Breeding Farm in Inner Mongolia Province, China. Twenty-one embryos (three samples at each stage) were randomly collected and any lineage was avoided during the sampling process. Skin samples (approximately 1 cm<sup>2</sup> for each individual) were collected from right mid-side of embryos at seven different embryonic days (E70, E75, E80, E90, E100, E120 and E130). Tissue was frozen in liquid nitrogen and stored at -80 °C

**Table 1** The RT-PCR and qRT-PCR primers used in this study

Primer Name	Sequence (5'-3')	Fragment size (bp)	T.M. (°C)
<i>Cloning primers</i>			
<i>TR<math>\alpha</math>-1 F</i>	CCTGGATGGAATTGAAGTGA	799	62.0
<i>TR<math>\alpha</math>-1R</i>	GACATGATCTCCATGCAGC		
<i>TR<math>\alpha</math>-2 F</i>	AGGCCTTCAGCGAGTTTAC	652	59.0
<i>TR<math>\alpha</math>-2R</i>	CCTTCTCTCCAGGCTCCTC		
<i>CRABPII-1 F</i>	CAGTGCTCCAGTGAAAGA	563	56.5
<i>CRABPII-1R</i>	CCAGAAGTGATTGGGTGAG		
<i>Real-time PCR primers</i>			
<i>TR<math>\alpha</math>-3 F</i>	TTACCTGGACAAAGACGAGC	113	57.4
<i>TR<math>\alpha</math>-3R</i>	TCTGGATTGTGCGGCGAAAG		
<i>CRABPII-2 F</i>	ACATCAAAACCTCCACCACC	111	56.5
<i>CRABPII-2R</i>	CCCATTTCACCAGGCTCTTA		
<i>ACTB-F</i>	CCTGCGGCATTCACGAACTAC	87	58.5
<i>ACTB-R</i>	ACAGCACCGTGTGGCGTAGAG		
<i>GAPDH-F</i>	GCA AGTTCCACGGCACAG	249	59.0
<i>GAPDH-R</i>	GGT TCACGCCCATCACAA		
<i>TOP2B-F</i>	GTGTGGAGCCTGAGTGGTATA	137	59.0
<i>TOP2B-R</i>	AAGCATTGCGCTGACATTGTT		


**(a)**

Mouse_NP_835161	MEQKPSKVECGSDPEENSARS PDGKRKRKNGQCSLKTSMISGYIPSYLDKDEQCVCVCGDKATGYHYRCITCEGCKGFFRRTIQKN	84
Rat_NP_001017960	MEQKPSKVECGSDPEENSARS PDGKRKRKNGQCSLKTSMISGYIPSYLDKDEQCVCVCGDKATGYHYRCITCEGCKGFFRRTIQKN	84
Human_NP_955366	MEQKPSKVECGSDPEENSARS PDGKRKRKNGQCSLKTSMISGYIPSYLDKDEQCVCVCGDKATGYHYRCITCEGCKGFFRRTIQKN	84
Pig_NP_999355	MEQKPSKVECGSDPEENSARS PDGKRKRKNGQCSLKTSMISGYIPSYLDKDEQCVCVCGDKATGYHYRCITCEGCKGFFRRTIQKN	84
Sheep_NP_001094389	MEQKPSKVECGSDPEENSARS PDGKRKRKNGQCSLKTSMISGYIPSYLDKDEQCVCVCGDKATGYHYRCITCEGCKGFFRRTIQKN	84
Goat_Tra	MEQKPSKVECGSDPEENSARS PDGKRKRKNGQCSLKTSMISGYIPSYLDKDEQCVCVCGDKATGYHYRCITCEGCKGFFRRTIQKN	84
Cattle_XP_005220791	MEQKPSKVECGSDPEENSARS PDGKRKRKNGQCSLKTSMISGYIPSYLDKDEQCVCVCGDKATGYHYRCITCEGCKGFFRRTIQKN	84
Chicken_NP_990644	MEQKPSKVECGSDPEENSARS PDGKRKRKNGQCSLKTSMISGYIPSYLDKDEQCVCVCGDKATGYHYRCITCEGCKGFFRRTIQKN	82
Consensus	meqkps s pe r dgkrkr kc qc k smsgyipsyldkdeqcvcvcdkatgyhyrcitcegckgffrriqkn	
Mouse_NP_835161	LHPTYSCKYDSCCVIDK ITRNQQLCRFKKCI AVGMAMDLVLDSSKRVAKRKLIEONRERRRKEEMIRSLQORPEPTPEEWDLI	168
Rat_NP_001017960	LHPTYSCKYDSCCVIDK ITRNQQLCRFKKCI AVGMAMDLVLDSSKRVAKRKLIEONRERRRKEEMIRSLQORPEPTPEEWDLI	168
Human_NP_955366	LHPTYSCKYDSCCVIDK ITRNQQLCRFKKCI AVGMAMDLVLDSSKRVAKRKLIEONRERRRKEEMIRSLQORPEPTPEEWDLI	168
Pig_NP_999355	LHPTYSCKYDSCCVIDK ITRNQQLCRFKKCI AVGMAMDLVLDSSKRVAKRKLIEONRERRRKEEMIRSLQORPEPTPEEWDLI	168
Sheep_NP_001094389	LHPTYSCKYDSCCVIDK ITRNQQLCRFKKCI AVGMAMDLVLDSSKRVAKRKLIEONRERRRKEEMIRSLQORPEPTPEEWDLI	168
Goat_Tra	LHPTYSCKYDSCCVIDK ITRNQQLCRFKKCI AVGMAMDLVLDSSKRVAKRKLIEONRERRRKEEMIRSLQORPEPTPEEWDLI	168
Cattle_XP_005220791	LHPTYSCKYDSCCVIDK ITRNQQLCRFKKCI AVGMAMDLVLDSSKRVAKRKLIEONRERRRKEEMIRSLQORPEPTPEEWDLI	168
Chicken_NP_990644	LHPTYSCKYDSCCVIDK ITRNQQLCRFKKCI AVGMAMDLVLDSSKRVAKRKLIEONRERRRKEEMIRSLQORPEPTPEEWDLI	166
Consensus	lhptysckyd ccvidkitrnqqlcrfkkci avgmamdlvldsskrvakrklie nrerrrkeemi slq rp p eew li	
Mouse_NP_835161	HVATEAHRSTNAQGS HWKQRKFLPDDIGQSP IIVSMPDGDVLDLEAFSEFTKIITPAITRVVDFAKKLPMSFELPCEDQIILLK	252
Rat_NP_001017960	HVATEAHRSTNAQGS HWKQRKFLPDDIGQSP IIVSMPDGDVLDLEAFSEFTKIITPAITRVVDFAKKLPMSFELPCEDQIILLK	252
Human_NP_955366	HVATEAHRSTNAQGS HWKQRKFLPDDIGQSP IIVSMPDGDVLDLEAFSEFTKIITPAITRVVDFAKKLPMSFELPCEDQIILLK	252
Pig_NP_999355	HVATEAHRSTNAQGS HWKQRKFLPDDIGQSP IIVSMPDGDVLDLEAFSEFTKIITPAITRVVDFAKKLPMSFELPCEDQIILLK	252
Sheep_NP_001094389	HVATEAHRSTNAQGS HWKQRKFLPDDIGQSP IIVSMPDGDVLDLEAFSEFTKIITPAITRVVDFAKKLPMSFELPCEDQIILLK	252
Goat_Tra	HVATEAHRSTNAQGS HWKQRKFLPDDIGQSP IIVSMPDGDVLDLEAFSEFTKIITPAITRVVDFAKKLPMSFELPCEDQIILLK	252
Cattle_XP_005220791	HVATEAHRSTNAQGS HWKQRKFLPDDIGQSP IIVSMPDGDVLDLEAFSEFTKIITPAITRVVDFAKKLPMSFELPCEDQIILLK	252
Chicken_NP_990644	HVATEAHRSTNAQGS HWKQRKFLPDDIGQSP IIVSMPDGDVLDLEAFSEFTKIITPAITRVVDFAKKLPMSFELPCEDQIILLK	250
Consensus	h teahrstnaqgshwkqrkflp digqsp iiv smpdgdvldleafseftkiitpaitrvvdfakklpmsfelpcedqiillk	
Mouse_NP_835161	GCCMEIMSLRAAVRYDPESD TLTLSGEMAVKREQLKNGGLGVVSDAIFELGKLSLAFNLDDTEVALLQAVLLMSTRSGLICVD	336
Rat_NP_001017960	GCCMEIMSLRAAVRYDPESD TLTLSGEMAVKREQLKNGGLGVVSDAIFELGKLSLAFNLDDTEVALLQAVLLMSTRSGLICVD	336
Human_NP_955366	GCCMEIMSLRAAVRYDPESD TLTLSGEMAVKREQLKNGGLGVVSDAIFELGKLSLAFNLDDTEVALLQAVLLMSTRSGLICVD	336
Pig_NP_999355	GCCMEIMSLRAAVRYDPESD TLTLSGEMAVKREQLKNGGLGVVSDAIFELGKLSLAFNLDDTEVALLQAVLLMSTRSGLICVD	336
Sheep_NP_001094389	GCCMEIMSLRAAVRYDPESD TLTLSGEMAVKREQLKNGGLGVVSDAIFELGKLSLAFNLDDTEVALLQAVLLMSTRSGLICVD	336
Goat_Tra	GCCMEIMSLRAAVRYDPESD TLTLSGEMAVKREQLKNGGLGVVSDAIFELGKLSLAFNLDDTEVALLQAVLLMSTRSGLICVD	336
Cattle_XP_005220791	GCCMEIMSLRAAVRYDPESD TLTLSGEMAVKREQLKNGGLGVVSDAIFELGKLSLAFNLDDTEVALLQAVLLMSTRSGLICVD	336
Chicken_NP_990644	GCCMEIMSLRAAVRYDPESD TLTLSGEMAVKREQLKNGGLGVVSDAIFELGKLSLAFNLDDTEVALLQAVLLMSTRSGLICVD	334
Consensus	gccmeimslraavrydpes tltlsgemavkreqlkngglgvvsdaif elgklsfafnlDDTEVALLQAVLLMSTRSGLICVD	
Mouse_NP_835161	KIEKSEQAYLLAFEHYVNRKHNIPHFWPKLLMKVTDLRMIGACHASRFLHMKVECPTELPFPLFLEVFEDQE	409
Rat_NP_001017960	KIEKSEQAYLLAFEHYVNRKHNIPHFWPKLLMKVTDLRMIGACHASRFLHMKVECPTELPFPLFLEVFEDQE	409
Human_NP_955366	KIEKSEQAYLLAFEHYVNRKHNIPHFWPKLLMKVTDLRMIGACHASRFLHMKVECPTELPFPLFLEVFEDQE	409
Pig_NP_999355	KIEKSEQAYLLAFEHYVNRKHNIPHFWPKLLMKVTDLRMIGACHASRFLHMKVECPTELPFPLFLEVFEDQE	409
Sheep_NP_001094389	KIEKSEQAYLLAFEHYVNRKHNIPHFWPKLLMKVTDLRMIGACHASRFLHMKVECPTELPFPLFLEVFEDQE	409
Goat_Tra	KIEKSEQAYLLAFEHYVNRKHNIPHFWPKLLMKVTDLRMIGACHASRFLHMKVECPTELPFPLFLEVFEDQE	409
Cattle_XP_005220791	KIEKSEQAYLLAFEHYVNRKHNIPHFWPKLLMKVTDLRMIGACHASRFLHMKVECPTELPFPLFLEVFEDQE	409
Chicken_NP_990644	KIEKSEQAYLLAFEHYVNRKHNIPHFWPKLLMKVTDLRMIGACHASRFLHMKVECPTELPFPLFLEVFEDQE	407
Consensus	kiek seqayllafehy n rkhnip hfwpkllmkvtdlrmigachas rflhmkvecptelfpplflevfedqe	

**(b)**

Sheep_XP_004002676	MPNFSGNWKIIRSENFEDLLKVLGNVMMRKAIVAAASKPAVEIKQEGD.TFYIKTSTTVRTEINFKVGEFEEQTVDRPCK	83
Goat_CRABPII	MPNFSGNWKIIRSENFEDLLKVLGNVMMRKAIVAAASKPAVEIKQEGD.TFYIKTSTTVRTEINFKVGEFEEQTVDRPCK	83
Pig_NP_001157981	MPNFSGNWKIIRSENFEDLLKVLGNVMMRKAIVAAASKPAVEIKQEGD.TFYIKTSTTVRTEINFKVGEFEEQTVDRPCK	83
Cattle_NP_001008670	MPNFSGNWKIIRSENFEDLLKVLGNVMMRKAIVAAASKPAVEIKQEGD.TFYIKTSTTVRTEINFKVGEFEEQTVDRPCK	83
Human_NP_001869	MPNFSGNWKIIRSENFEDLLKVLGNVMMRKAIVAAASKPAVEIKQEGD.TFYIKTSTTVRTEINFKVGEFEEQTVDRPCK	83
Mouse_NP_031785	MPNFSGNWKIIRSENFEDLLKVLGNVMMRKAIVAAASKPAVEIKQEND.TFYIKTSTTVRTEINFKVGEFEEQTVDRPCK	83
Rat_NP_058940	MPNFSGNWKIIRSENFEDLLKVLGNVMMRKAIVAAASKPAVEIKQEND.TFYIKTSTTVRTEINFKVGEFEEQTVDRPCK	84
Consensus	mpnfsngwkiirsenfe llk vlgvnm rkiavaaaskpaveikq d tfyiktsttvrtteinfk geefeeqtvdrpck	
Sheep_XP_004002676	SLVKWESENKMVCEQRLLKGEKPKTSWRELNDGELILMTADDIVCTRVYVR	137
Goat_CRABPII	SLVKWESENKMVCEQRLLKGEKPKTSWRELNDGELILMTADDIVCTRVYVR	137
Pig_NP_001157981	SLVKWESENKMVCEQRLLKGEKPKTSWRELNDGELILMTADDIVCTRVYVR	137
Cattle_NP_001008670	SLVKWESENKMVCEQRLLKGEKPKTSWRELNDGELILMTADDIVCTRVYVR	137
Human_NP_001869	SLVKWESENKMVCEQRLLKGEKPKTSWRELNDGELILMTADDIVCTRVYVR	137
Mouse_NP_031785	SLVKWESENKMVCEQRLLKGEKPKTSWRELNDGELILMTADDIVCTRVYVR	137
Rat_NP_058940	SLVKWESENKMVCEQRLLKGEKPKTSWRELNDGELILMTADDIVCTRVYVR	138
Consensus	slvkwes enkmvceqr llkgepktsw relndgelilmtaddivctrvyvr	

**Fig. 1** Alignment of the *Tra* (a) and *CRABPII* (b) amino acid sequences



**Fig. 2** The phylogenetic trees constructed by coding sequences of *TRα* (a) and *CRABP11* (b) based on the Neighbor-Joining method

for further analysis. All the experimental procedures for this experiment were conducted under a protocol approved by the Institutional Animal Care and Use Committee in the College of Animal Science and Technology, Sichuan Agricultural University, China.

#### RNA isolation and cDNA synthesis

The frozen skin tissues were ground using mortars in liquid nitrogen and the total RNA was isolated by Trizol reagent (Invitrogen, Carlsbad CA, USA) according to the manufacturer's protocols. The concentration and quality of the total RNA were further assessed using the Nano-Drop spectrophotometer (Bio-Rad, Benicia, USA). The RNase-free DNase I (Promega, Madison, USA) was used to digest genomic DNA. The first-strand cDNA was synthesized using the M-MLV reverse transcriptase kit (Promega, Madison, USA) with oligo (dT) primer.

#### Gene cloning and quantitative PCR analysis


Three primer pairs were designed to amplify the caprine *TRα* and *CRABP11* genes according to their conserved regions of homologies from human, mouse, cattle, sheep and pig (Table 1). PCR was carried out in a 25  $\mu$ L reaction mixture containing 2  $\mu$ L first-strand cDNA, 50 mM KCl, 10 mM Tris-HCl (pH 8.3), 0.5 mM MgCl<sub>2</sub>, 10 pmol each primer, 150  $\mu$ M dNTPs and 1 unit *Taq* polymerase (TaKaRa, Dalian, China). The cycling condition included an initial denaturation step at 95  $^{\circ}$ C for 5 min, 38 cycles of at 94  $^{\circ}$ C for 30 s, annealing temperature for 30 s and extension at 72  $^{\circ}$ C for 45 s, and a final extension at 72  $^{\circ}$ C for

7 min in a PTC-100 PCR thermocycler (MJ Research, Inc., Watertown, MA). PCR products were ligated with the pMD19-T vector (TaKaRa, Dalian, China) after purification, and sequenced by Invitrogen Biotech Co. Ltd. (Shanghai, China).

The quantitative PCR (qRT-PCR) was carried out in an iCycler iQ Real-Time PCR Detection System (Bio-Rad, Benicia, USA) with a total volume of 20  $\mu$ L containing 10  $\mu$ L 2 $\times$  SYBR Premix Ex Taq II, 0.6  $\mu$ L primers (10  $\mu$ M) and 1  $\mu$ L diluted cDNA. PCR reaction was as follows: a 95  $^{\circ}$ C denaturation for 30 s, followed by 40 cycles of 94  $^{\circ}$ C for 15 s, annealing temperature for 30 s, and 72  $^{\circ}$ C for 30 s. A melting program ranging from 55  $^{\circ}$ C to 95  $^{\circ}$ C with a heating rate of 0.5  $^{\circ}$ C/10 s was carried out to create the melt curves. Reactions were performed in triplicate and negative control was also performed in parallel.

#### Normalization of the expression data

In the present study, three internal control genes (*ACTB*, *GAPDH* and *TOP2B*, Table 1) were selected to normalize the expression levels of *TRα* and *CRABP11* mRNAs. To accurate expression profiling of target genes, the geometric mean of multiple carefully selected housekeeping genes was validated as an accurate normalization factor [24]. The relative gene expression was calculated with the  $2^{-\Delta\Delta C_t}$  method [25]. Data were presented as mean  $\pm$  SE. Comparisons between groups were analyzed *via* GLM (General Linear Model) for experiments with more than 2 subgroups. The significance level was  $P < 0.05$ .



**Fig. 3** The quantitative expressions of *TRα* (a) and *CRABP II* mRNAs (b) in skin tissue of Inner Mongolian Cashmere goat. The bar height presented the means, and error bar displayed +1SE (n = 3). Different letters above the bars indicate a significant difference (P < 0.05) between different stages

## Results and discussion

### Characteristics of goat *TRα* and *CRABP II* mRNAs

A 1,309-bp fragment of *TRα* was assembled by the two overlapped sequences of *TRα*-1 F/1R and *TRα*-2 F/2R with an open reading frame (ORF) extending from nucleotide positions 21 to 1,253 (with reference to the translational start codon of ATG), which encoded a protein with 410 amino acids (Accession No. KF589923). The obtained sequence of *CRABP II* mRNA was 563 bp in length with an ORF of 417 bp encoding 138 amino acids (Accession No. KF589924). The blast results revealed that both of *TRα* and *CRABP II* were quite conserved among species (Fig. 1, Additional file 1: Figure S1 and Additional file 2: Figure S2). The sequence similarity ranged from 88 % to 100 % (Additional file 3: Table S1). The coding sequence of the caprine *TRα* gene shows a high similarity with the sequences in other mammals, sharing 99 % identity with

sheep (NM\_001100919) and cattle (NM\_001046329). The goat *CRABP II* shows 88 % identity with mice (NM\_007759) and 98 % identity with cattle (NM\_001008670).

The nucleotide sequences were aligned by the Cluster W method included in the program BioEdit version 7.2.5 [26]. The phylogenetic analysis was constructed using the program MEGA 4.1 [27], with a Kimura 2-parameter model and a bootstrap test (1000 replications). The phylogenetic tree revealed that the goat *TR* grouped with sheep, and then clustered with cattle, pig, human, mice and chicken subsequently (Fig. 2a). The phylogenetic tree of *CRABP II* gene showed a similar clustering with differences in sort of branch-length groups (Fig. 2b). The Minimum Evolution, Maximum Parsimony and UPGMA trees revealed the same clustering groups as presented by the NJ trees (data not shown).

### Time-course expressions of *TRα* and *CRABP2* genes

To better understand the prenatal dynamical expressions of *TRα* and *CRABP2* in the skin tissue of cashmere goats, the qRT-PCR array was performed in the middle late embryonic stages (E70 to E130). As shown in Fig. 3, both of *TRα* and *CRABP2* mRNAs were detectable in all the tested time points. However, no significant difference of *TRα* gene expression was detected during the middle late development of goat embryos. The mRNA of *TRα* was expressed at E70 with relatively high level and mildly decreased in the following three stages (E75, E80, and E90), and then increased at E100 and reduced to the lowest level at E120, subsequently. The highest expression of *TRα* gene was observed in the last stage (E130,  $P > 0.05$ ). The previous studies has reported that the secondary follicles grew from E65 to E75 and then extended to skin surface. The complete structure of the secondary follicle was formed at E135 in Chinese cashmere goats [4, 28, 29]. Synchronously coupled with the early formation and growth of cashmere, the mRNA expression of *TRα* gene was up-regulated indicating that *TRα* could play a role in the time-course growth of goat cashmere.

The expression pattern of *CRABP2* mRNA showed an “up-down-up” trend, which revealed a significantly highest expression at E75 ( $P < 0.05$ ), and was down-regulated during E80 to E120 ( $P < 0.05$ ) then increased again at E130. In embryonic development of hair follicles, the glandula sebacea cells were observed in the skin tissue from cashmere goat fetus at E85 [30]. The glandula sebacea formed at E90 and accelerated the growth of primary hair follicles. However, the physiologic difference between primary and secondary follicles was that no glandula sebacea was found in secondary hair follicles. The second hair follicles grew retard and partially matured at E130. The mRNA expression of *CRABP2* at E90 was lower than that at E80 when no glandula sebacea was formed. The *CRABP2* gene could regulate the early development of glandula sebacea though modifying the concentration of RA. The mRNA of *CRABP2* gene at E100 expressed significantly higher than that at E120, which led more RA transported into nucleus and bound to its receptor, and proposed to boost the growth of glandula sebacea. In humans, the concentration of RA in cells could increase the mRNA expression of *CRABP2* in skin [31, 32].

In this study, we characterized the caprine *TRα* and *CRABP2* genes and quantified their mRNA expressions during the formation of secondary hair follicles in the middle late embryonic periods. Our study will enrich the knowledge of goat *TRα* and *CRABP2* genes and provide the foundation for further insight into their functions on cashmere growth.

### Conclusions

Cashmere wool is the very valuable production obtained from goats. It is very important to investigate the expressions of key functional genes associated with cashmere growth during the prenatal and process-oriented periods (from anagen to catagen and finally telogen). Taken together, our results profiled the expressions of *TRα* and *CRABP2* genes associated with prenatal development of goat hair follicle.

### Additional files

**Additional file 1: Figure S1.** Alignment of the *TRα* coding sequences in mammals (PPT 230 kb)

**Additional file 2: Figure S2.** Alignment of the *CRABP2* nucleotide sequences in mammals (PPT 146 kb)

**Additional file 3: Table S1.** The sequence similarity of *TRα* and *CRABP2* genes in this study (DOC 39 kb)

### Competing interests

The authors declare that they have no competing interests.

### Authors' contributions

HPZ and HL designed experiment, TZ, WZ and ZQZ conducted experiment, LL and LJW analyzed data, TZ and WZ written manuscript. All authors read and approved the final manuscript.

### Acknowledgements

This study was supported by the Chinese Domestic Animal Germplasm Resources Infrastructure, the Technology Support Program of Sichuan province (2011NZ0099-36) and the National Transgenic Project (2011ZX08008002).

Received: 12 April 2015 Accepted: 6 August 2015

Published online: 20 August 2015

### References

- Mou C, Jackson B, Schneider P, Overbeek PA, Headon DJ. Generation of the primary hair follicle pattern. *Proc Natl Acad Sci U S A*. 2006;103:9075–80.
- Cadau S, Rosignoli C, Rhetore S, Voegel J, Parenteau-Bareil R, Berthod F. Early stages of hair follicle development: a step by step microarray identity. *Eur J Dermatol*. 2013. doi:10.1684/ejd.2013.1972.
- Zhang JY, Yin J, Li JQ, Li CQ. Study on hair follicle structure and morphogenesis of the Inner Mongolian Arbas cashmere goat. *Sci Agric Sin*. 2007;40:1017–23.
- Li YR, Fan WB, Li CQ, Yin J, Zhang JY, Li JQ. Histomorphology research of the secondary follicle cycling of Inner Mongolia cashmere goat. *Sci Agric Sin*. 2008;41:3920–6.
- Li CQ, Yin J, Zhang HY, Guo ZC, Zhang WG, Gao AQ, et al. Comparative study on skin and hair follicles cycling between Inner Mongolia and Liaoning cashmere goats. *Acta Veterinariae Zootechnica Sinica*. 2005;36:674–9.
- Parry AL, Norton BW, Restall BJ. Skin follicle development in the Australian cashmere goat. *Aust J Agr Res*. 1992;43:857–70.
- Stenn KS, Paus R. Controls of hair follicle cycling. *Physiol Rev*. 2001;81:449–94.
- Jiang W, Yang YX, Xue P, Huang YJ, Chen YL. Identification of genes preferentially expressed in goat hair follicle anagen-catagen transition using suppression subtractive hybridization. *Anim Biotechnol*. 2012;23:11–23.
- Jin M, Wang L, Li S, Xing MX, Zhang X. Characterization and expression analysis of KAP7.1, KAP8.2 gene in Liaoning new-breeding cashmere goat hair follicle. *Mol Biol Rep*. 2011;38:3023–8.
- Wang X, Zhao ZD, Xu HR, Qu L, Zhao HB, Li T, et al. Variation and expression of KAP9.2 gene affecting cashmere trait in goats. *Mol Biol Rep*. 2012;39:10525–9.

11. Su R, Li JQ, Zhang WG, Yin J, Zhao J, Chang ZL. Expression of BMP2 in the Skin and Hair Follicle from Different Stage in Inner Mongolia Cashmere Goat. *Sci Agric Sin*. 2008;41:559–63.
12. Bai WL, Yin RH, Jiang WQ, Luo GB, Yin RL, Li C, et al. Molecular characterization of prolactin cDNA and its expression pattern in skin tissue of Liaoning Cashmere goat. *Biochem Genet*. 2012;50:694–701.
13. Seki Y, Yokohama M, Wada K, Fujita M, Kotani M, Nagura Y, et al. Expression analysis of the type I keratin protein keratin 33A in goat coat hair. *Anim Sci J*. 2011;82:773–81.
14. Shah RM, Ganai TA, Sheikh FD, Shanaz S, Shabir M, Khan HM. Characterization and polymorphism of keratin associated protein 1.4 gene in goats. *Gene*. 2013;518:431–42.
15. Safer JD. Thyroid hormone action on skin. *Curr Opin Endocrinol Diabetes Obes*. 2012;19:388–93.
16. van Beek N, Bodo E, Kromminga A, Gaspar E, Meyer K, Zmijewski MA, et al. Thyroid hormones directly alter human hair follicle functions: anagen prolongation and stimulation of both hair matrix keratinocyte proliferation and hair pigmentation. *J Clin Endocrinol Metab*. 2008;93:4381–8.
17. Villar D, Nicol F, Arthur JR, Dicks P. P, Cannavan A, Kennedy DG, et al. Type II and type III monodeiodinase activities in the skin of untreated and propylthiouracil-treated cashmere goats. *Res Vet Sci*. 2000;68:119–23.
18. Rhind SM, Kyle CE. Skin deiodinase profiles and associated patterns of hair follicle activity in cashmere goats of contrasting genotypes. *Aust J Agr Res*. 2004;55:443–8.
19. Torma H, Karlsson T, Michaelsson G, Rollman O, Vahlquist A. Decreased mRNA levels of retinoic acid receptor alpha, retinoid X receptor alpha and thyroid hormone receptor alpha in lesional psoriatic skin. *Acta Derm Venereol*. 2000;80:4–9.
20. Moeller LC, Cao X, Dumitrescu AM, Seo H, Refetoff S. Thyroid hormone mediated changes in gene expression can be initiated by cytosolic action of the thyroid hormone receptor  $\beta$  through the phosphatidylinositol 3-kinase pathway. *Nucl Recept Signal*. 2006;4, e020.
21. Everts HB, Sundberg JP, King Jr LE, Ong DE. Immunolocalization of enzymes, binding proteins, and receptors sufficient for retinoic acid synthesis and signaling during the hair cycle. *J Invest Dermatol*. 2007;127:1593–604.
22. Sessler RJ, Noy N. A ligand-activated nuclear localization signal in cellular retinoic acid binding protein-II. *Mol Cell*. 2005;18:343–53.
23. Everts HB, King Jr LE, Sundberg JP, Ong DE. Hair cycle-specific immunolocalization of retinoic acid synthesizing enzymes Aldh1a2 and Aldh1a3 indicate complex regulation. *J Invest Dermatol*. 2004;123:258–63.
24. Vandesompele J, De Preter K, Pattyn F, Poppe B, Van Roy N, De Paep A, et al. Accurate normalization of real-time quantitative RT-PCR data by geometric averaging of multiple internal control genes. *Genome Biol*. 2002;3:research0034.
25. Livak KJ, Schmittgen TD. Analysis of relative gene expression data using real-time quantitative PCR and the  $2^{-\Delta\Delta C(T)}$  Method. *Methods*. 2001;25:402–8.
26. Hall TA. BioEdit: a user-friendly biological sequence alignment editor and analysis program for Windows 95/98/NT. *Nucl Acids Symp Ser*. 1999;41:95–8.
27. Tamura K, Dudley J, Nei M, Kumar S. MEGA4: Molecular Evolutionary Genetics Analysis (MEGA) software version 4.0. *Mol Biol Evol*. 2007;24:1596–9.
28. Wang L, Peng L, Zhang W, Zhang J, Yang W, Ding L, et al. Initiation and development of skin follicles in the Inner Mongolian cashmere goat. *Acta Veterinariae Zootechnica Sinica*. 1996;27:524–30.
29. Zhang JX, Wang L, Yin J, Li JQ, Zhang HJ. Expression of KAP6 gene family on the skin of fetal goat. *Animal Husbandry and Feed Science*. 2009;30:20–1.
30. Zhang YJ, Li CQ, Li JQ. Study on Development of Skin and Hair Follicle from Fetal Inner Mongolian Arbas Cashmere Goats. *Acta Veterinariae Zootechnica Sinica*. 2006;37:761–8.
31. Astrom A, Tavakkol A, Pettersson U, Cromie M, Elder JT, Voorhees JJ. Molecular cloning of two human cellular retinoic acid-binding proteins (CRABP). Retinoic acid-induced expression of CRABP-II but not CRABP-I in adult human skin in vivo and in skin fibroblasts in vitro. *J Biol Chem*. 1991;266:17662–6.
32. Millar SE. Molecular mechanisms regulating hair follicle development. *J Invest Dermatol*. 2002;118:216–25.

**Submit your next manuscript to BioMed Central and take full advantage of:**

- Convenient online submission
- Thorough peer review
- No space constraints or color figure charges
- Immediate publication on acceptance
- Inclusion in PubMed, CAS, Scopus and Google Scholar
- Research which is freely available for redistribution

Submit your manuscript at  
[www.biomedcentral.com/submit](http://www.biomedcentral.com/submit)

